



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

NOTE ON THE BINARY STAR β 1212 = 24 AQUARI

From the date of its discovery by Burnham in 1890 until 1916 the angular motion in the fine double star β 1212 (R. A. $21^h34^m.4$; Decl. $-0^{\circ}30'$; Magnitudes 7.7, 8.2) was slow, the entire change in the 26 years amounting to only 40 degrees. The distance did not vary greatly from $0''.5$.

My measures in the present season (the only ones known to me since 1916), however, show a very marked change; the angular velocity has become much greater and the distance has diminished to less than half its earlier value. Burnham's conjecture, in 1899, that the components then had about their maximum separation seems to be confirmed.

The following observed positions will show the nature of the motion:

1890.75	$254^{\circ}.5$	0 .45	3n	Burnham
1898.80	$269^{\circ}.0$	0 .50	4	Burnham and Aitken
1916.52	$294^{\circ}.8$	0 .50	6	Leavenworth and Olivier
1921.66	$321^{\circ}.1$	0 .22	3	Aitken

R. G.AITKEN.

September 20, 1921.

A NEW DOUBLE STAR

On one of the plates taken in 1915 with the 30-inch refractor of the Allegheny Observatory for measuring the positions of the *Pleiades* stars the following B. D. star was found double which has not previously been listed as such:

B. D. $22^{\circ}536$, 9.3, R. A. $3^h37^m57^s.6$, Decl. $+22^{\circ}54'17''.5$ (1900)

The measures of the plate of discovery together with recent micro-metrical measures with the 12-inch refractor of the Lick Observatory give:

1915.98 Pos. A. $315^{\circ}.3$ Dist. $1''.56$ Photogr. images not well separated.

1921.64 $313^{\circ}.5$ I .55 3n.

The photographic magnitudes of the two components were estimated on the plate as 10.5 and 11.8; the spectral class of the *Henry Draper Catalogue* is A2. The star, tho situated in the *Pleiades*, is not a physical member of this group, but has a small proper-motion.

ROBERT TRUMPLER.